



Organic composition of carbonaceous aerosols in an aged prescribed fire plume

B. Yan, M. Zheng, Y. T. Hu, S. Yip Lee, H. K. Kim, A. G. Russell

► To cite this version:

B. Yan, M. Zheng, Y. T. Hu, S. Yip Lee, H. K. Kim, et al.. Organic composition of carbonaceous aerosols in an aged prescribed fire plume. *Atmospheric Chemistry and Physics Discussions*, 2007, 7 (6), pp.18015-18042. hal-00303215

HAL Id: hal-00303215

<https://hal.science/hal-00303215>

Submitted on 17 Dec 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Carbonaceous
aerosols in an aged
prescribed fire plume**

B. Yan et al.

Organic composition of carbonaceous aerosols in an aged prescribed fire plume

B. Yan¹, M. Zheng¹, Y. T. Hu², S. Lee¹, H. K. Kim², and A. G. Russell²

¹School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA

²School of Civil and Environmental Engineering, Georgia Institute of Technology, Atlanta, GA, USA

Received: 13 November 2007 – Accepted: 24 November 2007

– Published: 17 December 2007

Correspondence to: A. G. Russell (ted.russell@ce.gatech.edu)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Aged smoke from a prescribed fire (dominated by conifers) impacted Atlanta, GA on 28 February 2007 and dramatically increased hourly ambient concentrations of PM_{2.5} and organic carbon (OC) up to 140 and 72 $\mu\text{g m}^{-3}$, respectively. It was estimated that over 1 million residents were exposed to the smoky air lasting from the late afternoon to midnight. To better understand the processes impacting the aging of fire plumes, a detailed chemical speciation of carbonaceous aerosols was conducted by gas chromatography/mass spectrometry (GC/MS) analysis. Ambient concentrations of many organic species (levoglucosan, resin acids, retene, *n*-alkanes, *n*-alkanoic acids) associated with wood burning emission were significantly elevated on the event day. Levoglucosan increased by a factor of 10, while hopanes, steranes, cholesterol and major polycyclic aromatic hydrocarbons (PAHs) did not show obvious increases. Strong odd over even carbon number predominance was found for *n*-alkanes versus even over odd predominance for *n*-alkanoic acids. Alteration of resin acids during transport from burning sites to monitors is suggested by the observations. Our study also suggests that large quantities of biogenic volatile organic compounds (VOCs) and semivolatile organic compounds (SVOCs) were released both as products of combustion and unburned vegetation heated by the fire. Higher leaf temperature can stimulate biogenic VOC and SVOC emissions, which enhanced formation of secondary organic aerosols (SOA) in the atmosphere. This is supported by elevated ambient concentrations of secondary organic tracers (dicarboxylic acids, 2-methyltetrols, pinonic acid and pinic acid). An approximate source profile was built for the aged fire plume to help better understand evolution of wood smoke emission and can be used for source apportionment.

ACPD

7, 18015–18042, 2007

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

1 Introduction

Wildland fire (wild fire and prescribed burning) is estimated to contribute about 20% of total fine particulate matter (PM_{2.5}) emissions in the United States (EPA, 2000). In 2006, a total of 96 385 wildland fires were reported to burn 39 958 square kilometers, 125% above the 10-year average (NIFC, 2007). Among these forests, 11 010 square kilometers were treated with prescribed fires, which is 1 659 square kilometers above last year's total and is the second highest since 1998 (NIFC, 2007). Such large and increasing emission contributions are of concern to air quality managers, particularly in areas near or above the applicable air quality standards. For example, the prescribed fires on 28 February 2007 in Georgia and later the Georgia-Florida wildfires lasting from April through May severely impacted Atlanta, Georgia with thick wood smoke (Hu et al., 2007¹; Lee et al., 2007). During such events, hourly concentrations of PM_{2.5} increased by over 100 µg m⁻³.

Previous studies have shown that the major component of PM_{2.5} from forest burning events is organic carbon (OC), accounting for 30–70% of PM_{2.5} mass (Nopmongcol et al., 2007; Robinson et al., 2004; Ward et al., 2006). However, detailed data speciating OC in PM_{2.5} impacted by wildland fires is sparse, and even less data is available for aged plumes. During the 28 February 2007 event, OC reached 72 µg m⁻³ at 06:00 p.m. and contributed approximately 51% of the ambient PM_{2.5} in Atlanta, GA (Lee et al., 2007). This smoke event provides an opportunity to characterize OC in such carbonaceous aerosols, further understand processes impacting the aging of fire plumes, and estimate the composition of prescribed fire-derived PM_{2.5} for source apportionment studies.

Numerous organic compounds have been identified and quantified from ambi-

¹Hu, Y. T., Odman, M. T., Chang, M. E., Jackson, W., Lee, S., Edgerton, E. S., Baumann, K., and Russell, A. G.: Simulation of Air Quality Impacts from Prescribed Fire on an Urban Area: Analysis of the 28 February 2007, Environ. Sci. Technol., in review, 2007.

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ent PM_{2.5} samples including *n*-alkanes, hopanes, steranes, alkanolic acids, alkanedioic acids, PAHs, resin acids, and others (syringols, levoglucosan, cholesterol, 2-methyltetrols, etc.). Some compounds are reasonably unique tracers for certain sources and are widely used to track specific sources of carbonaceous aerosols. Similar sets of organic species have also been measured for source emissions. Together, they can be used to quantify source impacts on ambient PM_{2.5}.

Source impacts from biomass burning are usually traced through a few organic tracers including levoglucosan, resin acids, syringols and retene. As a pyrolysis product of cellulose in wood biopolymers, levoglucosan has been considered a particularly useful molecular marker of biomass burning (Simoneit et al., 1999). With its large emission abundance and reasonable thermal stability in the atmosphere, levoglucosan is frequently used to assess air quality impacts from biomass burning (Fraser and Lakshmanan, 2000; Schauer and Cass, 2000). Resin acids are thermal alteration products of coniferous wood resins and emitted exclusively from softwood burning (various pines, firs, etc.) (Rogge et al., 1998; Simoneit et al., 1993; Standley and Simoneit, 1994). In contrast, hardwood combustion produces much higher quantities of syringols (Hawthorne et al., 1988; Hawthorne et al., 1989). Although PAHs are emitted from multiple combustion processes of fuels (biomass, natural gas, diesel and gasoline) and ubiquitous in the atmosphere, retene, a thermal alteration of abietane compounds (resin diterpenoids), is considered as an organic tracer specific for coniferous wood burning (Ramdahl, 1983).

Other primary sources can be also linked to some specific organic tracers. Hopanes and steranes are emitted from both gasoline-powered vehicle and diesel-powered vehicle (Simoneit, 1985; Zielinska et al., 2004). They have been widely used as molecular markers of vehicular emissions in source apportionment of PM_{2.5} and OC (Fraser et al., 2003b; Schauer et al., 1996; Zheng et al., 2002, 2006, 2007). Cholesterol, found in animal fats and oils, is thought as an excellent molecular marker of meat cooking emission (Rogge et al., 1991). Vegetative detritus emissions are characterized by high-molecular weight *n*-alkanes with pronounced odd over even carbon number

**Carbonaceous
aerosols in an aged
prescribed fire plume**

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

predominance (Rogge et al., 1993a).

In addition to primary components of PM_{2.5}, secondary organic aerosol (SOA) formation can result from gaseous emissions of isoprenoids (isoprene and monoterpene)(Claeys et al., 2004a; Kavouras et al., 1998). A few biogenic SOA species have been identified and quantified including 2-methyltetrols (oxidation products of isoprene), pinonic acid and pinic acid (oxidation products of monoterpene) (Claeys et al., 2004a, b; Yu et al., 1999a, b). Although dicarboxylic acids (alkanedioic acids and dicarboxylic aromatic acids) can be emitted from various primary sources (mobile emission, meat cooking, etc.), previous studies suggested that atmospheric photochemical formation is probably the main source of these dicarboxylic acids (Fine et al., 2004b; Fraser et al., 2003a; Schauer et al., 2002).

Here, we collected PM_{2.5} filter samples before, during and after the 28 February 2007 prescribed fire episode impacting Atlanta, GA and analyzed organic composition of carbonaceous aerosols using GC/MS. We further capitalize on the quantified organic tracers to better understand the evolution of wood smoke from wildland fires.

2 Method description

2.1 Ambient sampling

Daily PM_{2.5} samples were collected on 47 mm quartz fiber filters with particulate composition monitors (PCM) at the Assessment of Spatial Aerosol Composition in Atlanta (ASACA) sites (Butler et al., 2003). These sites are located in the Atlanta metro area about 80 km downwind from the 28 February 2007 prescribed fires and were impacted directly by the smoke plume. The 24-h daily samples were analyzed for organic carbon/elemental carbon (OC/EC), metals and ions using thermal optical transmittance (TOT), X-ray fluorescence (XRF) and ion chromatography (IC), respectively. Details of ambient sampling, chemical measurements and particle compositions are described elsewhere (Baumann et al., 2003; Lee et al., 2005, 2007).

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2.2 Organic speciation

Due to the limitation of low air volumes sampled, the OC mass on a single PCM filter is usually not enough for organic tracer analysis. Therefore, three composite PCM samples (named “Before_Fire”, “Event” and “After_Fire”) were prepared as below. “Before_Fire” was composed of 6 PCM filters from three ASACA sites for the two days before the smoke day. “Event” was composed of 3 PCM filters collected at the same sites on 28 February, the day most directly impacted. “After_Fire” was composed of 7 PCM filters from the same sites within the three days after the smoke day. The three composite samples along with a composite field blank were analyzed for organic compounds in $PM_{2.5}$ using a standardized method described elsewhere (Nolte et al., 2002; Zheng et al., 2002, 2006). Briefly, each filter composite was spiked with deuterated internal standard (IS) mixtures and then successively extracted using hexane and benzene/isopropanol (2:1, v/v). After filtering, extracts were concentrated with rotary evaporation followed by blowdown under pure nitrogen. Half of each concentrated extract was then derivatized with diazomethane to convert organic acids to their methyl esters. These methylated extracts were analyzed by GC/MS along with authentic standards. To quantify a few polar organic compounds (levoglucosan, cholesterol, 2-methyltetrols), underivatized remains of concentrated extracts were silylated with BSTFA (N, O-bis(trimethylsilyl)acetamide) to convert polar compounds to trimethylsilyl (TMS) derivatives. After one hour reaction under $70^{\circ}C$, these silylated extracts were analyzed using GC/MS along with authentic standards.

3 Results and discussion

Our results show that the observed 24-h average OC concentration jumped from 3.5 to $17.7 \mu g m^{-3}$ on the event day, accounting for more than 60% (when converted to organic matter by a factor of 1.4) of the total 24-h average $PM_{2.5}$ mass of $37 \mu g m^{-3}$. Major sources of carbonaceous aerosols are traced using variations of their associated

ACPD

7, 18015–18042, 2007

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

organic tracers.

3.1 Organic tracers of biomass burning

During the smoke, large increases were observed for biomass burning tracers (levoglucosan, resin acids, retene, etc.). Levoglucosan was detected in all ambient samples as the most abundant organic compound (Fig. 1). Before and after the smoke event, the observed levoglucosan concentrations were 114 and 145 ng m⁻³, respectively. On the event day, the levoglucosan concentration increased dramatically to 1210 ng m⁻³ and contributed 7% of the total OC, suggesting that the wood burning emission impact was 10 times higher on the event day than the non-smoke days. Along with levoglucosan, concentrations of resin acids also increased, especially dehydroabietic acid and 7-oxodehydroabietic acid, increasing to 42 and 19 ng m⁻³, approximately 9 and 23 times higher respectively than the levels before the burning day. Unlike levoglucosan, their ambient concentrations remained elevated after the fires (Fig. 2). To elucidate the processes occurring during transport, comparisons between wood burning source emissions and ambient data are conducted here. Ratios of major resin acids to levoglucosan were calculated and compared for both ambient data and a few source emissions from prominent softwood species in the southern United States (Table 1). In the softwood source emissions, abietic acid and dehydroabietic acid generally account for the majority of resin acids, about 58% and 32% respectively on average, while 7-oxodehydroabietic acid is minor. However, dehydroabietic acid and 7-oxodehydroabietic acid constituted a major fraction of the observed resin acids, about 65% and 29% respectively, in the ambient sample impacted by the fires. This comparison provides evidence that dehydroabietic acid and 7-oxodehydroabietic acid are being formed from other resin acids (i.e. diterpenoids) during the 3–4 h (around 80 km travel distance) transport from the burning sites to the monitors. Note that further alterations of dehydroabietic acid and 7-oxodehydroabietic acid exist, leading to their ratios to levoglucosan being variable over time during transport. Previous studies have also proposed that dehydroabietic acid and 7-oxodehydroabietic acid can be formed through

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

oxidation processes of other resin acids (i.e. abietic acids), resulting in accumulated concentrations in the atmosphere (Oros and Simoneit, 2001; Rogge et al., 1993b). Although most PAHs did not show significant elevation in ambient concentrations during the event day, retene, a potential softwood burning tracer, increased by around 12 and 7 times, respectively compared to before and after the fire, and predominated among all PAHs (Fig. 2).

n-alkanes and *n*-alkanoic acids identified in this study ranged from C₁₇ to C₃₆ and from C₁₄ to C₃₀, respectively (Fig. 3). Generally, *n*-alkanes are associated with plant wax and fossil fuel contributions, depending on carbon numbers. To approximately compare source impacts from plant wax versus fossil fuel combustion, the carbon preference index (CPI) measuring carbon number predominance in homologous compound series is calculated (Mazurek and Simoneit, 1984; Oros et al., 2006). In the samples before and after the fire, *n*-alkanes only show a slight odd carbon number predominance with CPIs of 1.4 and 1.7, respectively. However, *n*-alkanes exhibited distinctly strong odd carbon number predominance (CPI=3.1, carbon number maximum C_{max}=29) on the event day, reflecting a major contribution from plant waxes. Likewise, *n*-alkanoic acids had a strong even carbon number predominance on the event day (CPI=5.7, C_{max}=24). The concentration of even-over-odd carbon number series on the event day was 50 ng m⁻³, much larger than those before and after the event (12 and 10 ng m⁻³, respectively). Thus, a dominant contribution from plant waxes is suggested. These compounds are considered natural products of epicuticular waxes and internal lipid substances in leaf surfaces and emitted as vegetative detritus or through direct thermal volatilization (Rogge et al., 1993a; Simoneit, 2002). Higher leaf temperatures presumably lead to volatilization of these high-molecular weight organic compounds, which can then condense into the particle phase in the atmosphere. This result indicates that such waxes are not unique tracers for vegetative detritus when wildland fires also significantly impact air quality.

**Carbonaceous
aerosols in an aged
prescribed fire plume**

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.2 Other primary organic tracers

A few non-biomass organic molecular markers were also analyzed as indicators of other major primary sources. Vehicular engine exhaust and meat cooking are the two other major sources of primary organic carbon in urban areas. The before, during and after concentrations of hopanes and steranes did not show significant variation, i.e. 1.7, 1.7 and 1.0 ng m⁻³, indicating relatively consistent source contributions from vehicular emissions in the Atlanta area (Fig. 4). Cholesterol, an organic tracer of meat cooking, did not exhibit major variation as well (Fig. 5). Thus, it is clear that vehicular sources and meat cooking emissions are not responsible for the large increase in OC, which accounts for 81% of the total OC, on the event day, and that the greatly increased carbonaceous aerosol concentrations during the smoke episode are from prescribed fires. The source apportionment results calculated using organic molecular marker-based chemical mass balance (CMB-MM) model also indicate significant increases of prescribed burning emissions on the event day, but not for other major primary sources (i.e. vehicular source and meat cooking) (Lee et al., 2007). In addition, about 43% of the total observed OC can not be explained by primary source contributions, suggesting that secondary organic aerosol (SOA) might be a main contributor to the increased OC.

3.3 Secondary organic tracers

Our results for secondary organic tracers indicate that additional biogenic SOA were formed during the 28 February 2007 fire episode. 2-methyltetrols, secondary organic products of isoprene, were detected only on the smoke plume day with a concentration of 0.8 ng m⁻³ (Fig. 5). Pinonic acid and pinic acid, photo-oxidation products of monoterpenes, increased from 2.9 and 1.2 ng m⁻³ to 5.0 and 3.1 ng m⁻³, respectively, from the day prior to the fire through the smoke day (Fig. 5). Additionally, elevated dicarboxylic acids (alkanedioic acids and dicarboxylic aromatic acids) provided supportive evidence of increased SOA formation in the atmosphere. They are further

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

considered here products of biogenic emissions enhanced by the fire since significant increases were not found in anthropogenic emissions (i.e. vehicular source and meat cooking). The sums of dicarboxylic acids were 7.5, 51, 8.4 ng m⁻³ before, during and after the event, respectively. There are three possible hypotheses relating to the large increases in these dicarboxylic acids. The most direct one is that precursors of these compounds are associated with prescribed fire emissions. As stated previously, emissions of biogenic VOC and SVOC (isoprenes, terpenes, sesquiterpenes, alcohols, esters, carbonyls, acids, etc.) would be enhanced by either wood combustion process or increasing leaf temperature. Emissions of isoprenoids (isoprene and monoterpene) have been observed to be higher during forest fires due to increased temperatures (Alessio et al., 2004). A previous study also suggests that leaf temperatures rising from 25 to 35°C would increase biogenic VOC and SVOC emissions by 4 and 1.5 times from isoprene-emitting deciduous trees and terpene-emitting conifers, respectively (Lamb et al., 1987). The second hypothesis is that the dramatic increase in OC mass can enhance formation of SOA due to increased partitioning of these diacids to the particle phase. However, unless there is a significant chemical affinity between OC and the compounds, such a shift is not expected to be so large (less than 10% of enhancement in SOA) (Nopmongcol et al., 2007; Odum et al., 1996). The third hypothesis is that enhanced photo-oxidation occurred. The observed increase of ozone in the plume is indicative that this effect might exist, but would be limited since ozone increased about 50% (from around 60 to 90 ppb), significantly less than the increase in dicarboxylic acids (8 times higher). Therefore, among the three hypotheses above, the evidence strongly supports that biomass burning is a major contributor to precursors of dicarboxylic acids. Results of air quality modeling and measurement of water soluble organic carbon (WSOC) provide further supporting evidence (Hu et al., 2007¹; Lee et al., 2007).

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.4 Source profiles for an aged plume

Capturing the fire event provides information to assess the source composition profiles of an aged prescribed fire plume. Impacted by the fire plume, OC increased by $14.2 \mu\text{g m}^{-3}$, accounting for 81% of total observed OC on the event day while $\text{PM}_{2.5}$ increased by $23.9 \mu\text{g m}^{-3}$, accounting for 64% of total observed $\text{PM}_{2.5}$. Organic markers of other major primary emissions (vehicular source and meat cooking) did not show significant variation in Atlanta, GA during the episode. Thus, the large increases in OC, $\text{PM}_{2.5}$ and associated chemical species on the event day are considered a direct contribution from prescribed fire emissions. Here, two approximate source composition profiles were developed for the aged biomass burning plume by considering differences between the before_event day (non-fire event impacted) and the event day, designated “aged_plume profile” and “primary_plume profile”. “aged_plume profile” is constituted by fractions of individual chemical species (increased concentrations) in the total increased (fire-caused total) $\text{PM}_{2.5}$ mass, i.e. $23.9 \mu\text{g m}^{-3}$, on the event day (Table 2). This profile contains chemical compositions of the aged plume, but is not the primary $\text{PM}_{2.5}$ source profile due to large quantities of SOA in aerosol from the fire emissions. “primary_plume profile” is then calculated using fractions of individual chemical species (increased concentrations) in the fire-caused primary $\text{PM}_{2.5}$ mass where the estimated fire-caused secondary organic carbon (SOC) was subtracted from the fire-caused total $\text{PM}_{2.5}$ mass on the event day (Table 2). The fire-caused SOC on the event was estimated based on the CMB-MM modeling results and measured EC/OC ratios. Briefly, the CMB-MM apportionment results indicate that prescribed fires contributed approximately 1.1 and $8.7 \mu\text{g m}^{-3}$ to the total primary OC before and on the event day, respectively (Lee et al., 2007). The difference, $7.6 \mu\text{g m}^{-3}$, is attributed to the plumes. Results here find that the total, primary plus secondary, impact of the plumes on OC is $14.2 \mu\text{g m}^{-3}$, suggesting that $6.6 \mu\text{g m}^{-3}$ (47% of the fire-caused total OC) comes from enhanced SOC. A comparable SOC fraction in the fire-caused total OC was also estimated using the EC/OC ratio method. In the aged plume, the fire-caused EC/OC ratio

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

is 0.039 while a value of 0.065 has been measured for prescribed burning emissions in Georgia (Lee et al., 2005). The lower EC/OC ratio suggests SOA formation in the aged plume. Using those ratios, SOC is calculated to account for 40% of the fire-caused total OC. Averaging the two, 44% of fire-caused total OC is taken as SOC during the event, and the fire-caused primary OC (POC_{fc}) is then estimated by:

$$POC_{fc} = OC \text{ (fire - caused)} \times (1 - f_{SOC}) = [OC \text{ (event)} - OC \text{ (before - event)}] \times (1 - f_{SOC}), \quad (1)$$

where f_{SOC} is the estimated SOC fraction in the fire-caused total OC. Similarly, the amount of fire-caused primary $PM_{2.5}$ (PPM_{fc}) is calculated as:

$$PPM_{fc} = [PM \text{ (event)} - PM \text{ (before - event)}] - OC \text{ (fire - caused)} \times f_{SOC}. \quad (2)$$

Finally, fractions of individual species (f_i) in fire-caused primary $PM_{2.5}$ are found as:

$$f_i = \frac{C_i \text{ (event)} - C_i \text{ (before - event)}}{PPM_{fc}} \quad (3)$$

where C_i is the ambient concentration of individual chemical species in $PM_{2.5}$ that are viewed as being dominated by primary emissions. In addition, an overall uncertainty of each chemical species in the two plume source profiles was calculated by propagating uncertainties associated with the observed OC, estimated SOC fraction and measured chemical species (i.e. organic compounds, EC, ions and trace metals).

Our study indicates that the “primary plume profile” derived from the fire event is comparable with the prescribed burning emission profile measured by Lee et al. (2005) for some organic compounds, but significantly different for others (Table 2 and Fig. 6). In the aged fire plume, levoglucosan accounts for 14% of primary OC, comparable to 9.5% in the profile of Lee et al. (2005). Most *n*-alkanes also show comparable levels for the two source profiles. Hopanes and steranes were not significant in either set. The ratio of water-soluble potassium (K^+)/OC in the “primary plume profile” is 0.0067, lower than the values in Lee et al. (0.011) and in Fine et al. (0.012 on average), but comparable to the average ratio of 0.0058 from foliar fuel combustion (Fine et al., 2004a; Hays et

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

al., 2002; Lee et al., 2005). However, significant differences were found for the fractions of PAHs, resin acids, *n*-alkanoic acid and dicarboxylic acids, suggesting aging of fire smoke after a 3–4 h travel distance. Lee et al. (2005) measured higher abundances for many PAHs and resin acids, implying direct combustion is the main contributor of these compounds and significant chemical alteration occurs during transport, especially for resin acids. In contrast, the aged plume has distinctly higher dicarboxylic acids (alkanedioic acids and dicarboxylic aromatic acids) fractions. Enhanced SOA formation in the atmosphere from the fire is suggested by these secondary indicators. The profile comparison between prescribed fire source emission and aged plume is useful to understand evolution of wood smoke during transport and to assess the organic species applied in CMB-MM modeling. Impacts from aging process of smoke plume should be considered in source apportionment methodology using CMB model, i.e. selection of fitting species.

4 Conclusions

Large increases in PM_{2.5} and OC in Atlanta caused by an aged prescribed fire plume through direct burning emissions and formation of secondary organic aerosols was captured by a system of monitors. Organic tracers of biomass burning, levoglucosan, resin acids and retene, exhibited sharp increases in concentrations during the episode. Observed resin acids and retene indicated predominant softwood (conifers) burning. Increases in resin acids were accompanied by high levels of dehydroabietic acids and 7-oxodehydroabietic acids, mainly formed during transport. Carbon number predominance for *n*-alkanes (odd over even) and *n*-alkanoic acids (even over odd) suggested that emissions from heat-exposed vegetation are important as one of major sources for increasing OC, especially secondary OC. Secondary organic tracers were observed with significantly higher ambient concentrations, providing further support for biogenic SOA formation. These data allows estimating source profiles for aged fire plumes, which can be used for improving source apportionment

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of wood burning by CMB modeling <http://www.atmos-chem-phys-discuss.net/7/18015/2007/acpd-7-18015-2007-supplement.pdf>.

Acknowledgements. This work was funded by the U.S. Environmental Protection Agency STAR grants (R832159, R828976 and R831076). We would like to thank Georgia Power (Southern Company) for their support of work in Laboratory for Atmospheric Modeling, Diagnostics, and Analysis (LAMDA) at Georgia Institute of Technology in this study area. The authors also acknowledge the helpful forest information provided by D. Tian.

References

- Alessio, G. A., De Lillis, M., Fanelli, M., Pinelli, P., and Loreto, F.: Direct and indirect impacts of fire on isoprenoid emissions from Mediterranean vegetation, *Funct. Ecol.*, 18, 357–364, 2004.
- Baumann, K., Ift, F., Zhao, J. Z., and Chameides, W. L.: Discrete measurements of reactive gases and fine particle mass and composition during the 1999 Atlanta Supersite Experiment, *J. Geophys. Res.-Atmos.*, 108, 8416, doi:10.1029/2001jd001210, 2003.
- Butler, A. J., Andrew, M. S., and Russell, A. G.: Daily sampling of PM_{2.5} in Atlanta: results of the first year of the assessment of spatial aerosol composition in Atlanta study, *J. Geophys. Res.-Atmos.*, 108, 8415, doi:10.1029/2002jd002234, 2003.
- Claeys, M., Graham, B., Vas, G., Wang, W., Vermeylen, R., Pashynska, V., Cafmeyer, J., Guyon, P., Andreae, M. O., Artaxo, P., and Maenhaut, W.: Formation of secondary organic aerosols through photooxidation of isoprene, *Science*, 303, 1173–1176, 2004a.
- Claeys, M., Wang, W., Ion, A. C., Kourtshev, I., Gelencser, A., and Maenhaut, W.: Formation of secondary organic aerosols from isoprene and its gas-phase oxidation products through reaction with hydrogen peroxide, *Atmos. Environ.*, 38, 4093–4098, doi:10.1016/j.atmosenv.2004.06.001, 2004b.
- EPA, U. S.: National Air Pollutant Emission Trends 1900–1998, EPA 454/R-400–002, Office Washington, DC, 2000.
- Fine, P. M., Cass, G. R., and Simoneit, B. R. T.: Chemical characterization of fine particle emissions from fireplace combustion of woods grown in the northeastern United States, *Environ. Sci. Technol.*, 35, 2665–2675, 2001.

ACPD

7, 18015–18042, 2007

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

- Fine, P. M., Cass, G. R., and Simoneit, B. R. T.: Chemical characterization of fine particle emissions from the fireplace combustion of woods grown in the southern United States, *Environ. Sci. Technol.*, 36, 1442–1451, 2002.
- 5 Fine, P. M., Cass, G. R., and Simoneit, B. R. T.: Chemical characterization of fine particle emissions from the wood stove combustion of prevalent United States tree species, *Environ. Eng. Sci.*, 21, 705–721, 2004a.
- Fine, P. M., Chakrabarti, B., Krudysz, M., Schauer, J. J., and Sioutas, C.: Diurnal variations of individual organic compound constituents of ultrafine and accumulation mode particulate matter in the Los Angeles basin, *Environ. Sci. Technol.*, 38, 1296–1304, 2004b.
- 10 Fraser, M. P. and Lakshmanan, K.: Using levoglucosan as a molecular marker for the long-range transport of biomass combustion aerosols, *Environ. Sci. Technol.*, 34, 4560–4564, 2000.
- Fraser, M. P., Cass, G. R., and Simoneit, B. R. T.: Air quality model evaluation data for organics. 6. C-3-C-24 organic acids, *Environ. Sci. Technol.*, 37, 446–453, 2003a.
- 15 Fraser, M. P., Yue, Z. W., and Buzcu, B.: Source apportionment of fine particulate matter in Houston, TX, using organic molecular markers, *Atmos. Environ.*, 37, 2117–2123, 2003b.
- Hawthorne, S. B., Miller, D. J., Barkley, R. M., and Krieger, M. S.: Identification of Methoxylated Phenols as Candidate Tracers for Atmospheric Wood Smoke Pollution, *Environ. Sci. Technol.*, 22, 1191–1196, 1988.
- 20 Hawthorne, S. B., Krieger, M. S., Miller, D. J., and Mathiason, M. B.: Collection and Quantitation of Methoxylated Phenol Tracers for Atmospheric-Pollution from Residential Wood Stoves, *Environ. Sci. Technol.*, 23, 470–475, 1989.
- Hays, M. D., Geron, C. D., Linna, K. J., Smith, N. D., and Schauer, J. J.: Speciation of gas-phase and fine particle emissions from burning of foliar fuels, *Environ. Sci. Technol.*, 36, 2281–2295, 2002.
- 25 Kavouras, I. G., Mihalopoulos, N., and Stephanou, E. G.: Formation of atmospheric particles from organic acids produced by forests, *Nature*, 395, 683–686, 1998.
- Lamb, B., Guenther, A., Gay, D., and Westberg, H.: A National Inventory of Biogenic Hydrocarbon Emissions, *Atmos. Environ.*, 21, 1695–1705, 1987.
- 30 Lee, S., Baumann, K., Schauer, J. J., Sheesley, R. J., Naeher, L. P., Meinardi, S., Blake, D. R., Edgerton, E. S., Russell, A. G., and Clements, M.: Gaseous and particulate emissions from prescribed burning in Georgia, *Environ. Sci. Technol.*, 39, 9049–9056, doi:10.1021/Es0515831, 2005.

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Lee, S., Kim, H. K., Yan, B., Cobb, E., Hennigan, C., Nichols, S., Chamber, M., Edgerton, E. S., Jansen, J. J., Hu, Y. T., Zheng, M., Weber, R. J., and Russell, A. G.: Diagnosis of Aged Prescribed Fire Plumes Hitting an Urban Area: Implications to Air Quality Modeling, Environ. Sci. Technol., in press, 2007.

5 NIFC (National Interagency Fire Center): fire Information – wildland fire statistics, available at: http://www.nifc.gov/fire_info/fire_stats.htm, access: August, 2007.

Nolte, C. G., Schauer, J. J., Cass, G. R., and Simoneit, B. R. T.: Trimethylsilyl derivatives of organic compounds in source samples and in atmospheric fine particulate matter, Environ. Sci. Technol., 36, 4273–4281, 2002.

10 Nopmongkol, U., Khamwichit, W., Fraser, M. P., and Allen, D. T.: Estimates of heterogeneous formation of secondary organic aerosol during a wood smoke episode in Houston, Texas, Atmos. Environ., 41, 3057–3070, 2007.

Odum, J. R., Hoffmann, T., Bowman, F., Collins, D., Flagan, R. C., and Seinfeld, J. H.: Gas/particle partitioning and secondary organic aerosol yields, Environ. Sci. Technol., 30, 2580–2585, 1996.

15 Oros, D. R., and Simoneit, B. R. T.: Identification and emission factors of molecular tracers in organic aerosols from biomass burning Part 1. Temperate climate conifers, Appl. Geochem., 16, 1513–1544, 2001.

Oros, D. R., Bin Abas, M. R., Omar, N. Y. M. J., Rahman, N. A., and Simoneit, B. R. T.: Identification and emission factors of molecular tracers in organic aerosols from biomass burning: Part 3. Grasses, Appl. Geochem., 21, 919–940, 2006.

20 Ramdahl, T.: Retene – a Molecular Marker of Wood Combustion in Ambient Air, Nature, 306, 580–583, 1983.

Robinson, M. S., Chavez, J., Velazquez, S., and Jayanty, R. K. M.: Chemical speciation of PM_{2.5} fires of the coconino national collected during prescribed forest near Flagstaff, Arizona, J. Air Waste Manage., 54, 1112–1123, 2004.

25 Rogge, W. F., Hildemann, L. M., Mazurek, M. A., Cass, G. R., and Simoneit, B. R. T.: Sources of Fine Organic Aerosol .1. Charbroilers and Meat Cooking Operations, Environ. Sci. Technol., 25, 1112–1125, 1991.

30 Rogge, W. F., Hildemann, L. M., Mazurek, M. A., Cass, G. R., and Simoneit, B. R. T.: Sources of Fine Organic Aerosol .4. Particulate Abrasion Products from Leaf Surfaces of Urban Plants, Environ. Sci. Technol., 27, 2700–2711, 1993a.

Rogge, W. F., Mazurek, M. A., Hildemann, L. M., Cass, G. R., and Simoneit, B. R. T.: Quan-

**Carbonaceous
aerosols in an aged
prescribed fire plume**

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- tification of Urban Organic Aerosols at a Molecular-Level - Identification, Abundance and Seasonal-Variation, *Atmos. Environ. A-Gen.*, 27, 1309–1330, 1993b.
- Rogge, W. F., Hildemann, L. M., Mazurek, M. A., Cass, G. R., and Simoneit, B. R. T.: Sources of fine organic aerosol. 9. Pine, oak and synthetic log combustion in residential fireplaces, *Environ. Sci. Technol.*, 32, 13–22, 1998.
- Schauer, J. J., Rogge, W. F., Hildemann, L. M., Mazurek, M. A., and Cass, G. R.: Source apportionment of airborne particulate matter using organic compounds as tracers, *Atmos. Environ.*, 30, 3837–3855, 1996.
- Schauer, J. J. and Cass, G. R.: Source apportionment of wintertime gas-phase and particle-phase air pollutants using organic compounds as tracers, *Environ. Sci. Technol.*, 34, 1821–1832, 2000.
- Schauer, J. J., Fraser, M. P., Cass, G. R., and Simoneit, B. R. T.: Source reconciliation of atmospheric gas-phase and particle-phase pollutants during a severe photochemical smog episode, *Environ. Sci. Technol.*, 36, 3806–3814, 2002.
- Simoneit, B. R. T.: Application of Molecular Marker Analysis to Vehicular Exhaust for Source Reconciliations, *Int J. Environ. An. Ch.*, 22, 203–233, 1985.
- Simoneit, B. R. T., Rogge, W. F., Mazurek, M. A., Standley, L. J., Hildemann, L. M., and Cass, G. R.: Lignin Pyrolysis Products, Lignans, and Resin Acids as Specific Tracers of Plant Classes in Emissions from Biomass Combustion, *Environ. Sci. Technol.*, 27, 2533–2541, 1993.
- Simoneit, B. R. T., Schauer, J. J., Nolte, C. G., Oros, D. R., Elias, V. O., Fraser, M. P., Rogge, W. F., and Cass, G. R.: Levoglucosan, a tracer for cellulose in biomass burning and atmospheric particles, *Atmos. Environ.*, 33, 173–182, 1999.
- Simoneit, B. R. T.: Biomass burning - A review of organic tracers for smoke from incomplete combustion, *Appl. Geochem.*, 17, 129–162, 2002.
- Standley, L. J. and Simoneit, B. R. T.: Resin Diterpenoids as Tracers for Biomass Combustion Aerosols, *J. Atmos. Chem.*, 18, 1–15, 1994.
- Ward, T. J., Hamilton, R. F., Dixon, R. W., Paulsen, M., and Simpson, C. D.: Characterization and evaluation of smoke tracers in PM: Results from the 2003 Montana wildfire season, *Atmos. Environ.*, 40, 7005–7017, 2006.
- Yu, J. Z., Cocker, D. R., Griffin, R. J., Flagan, R. C., and Seinfeld, J. H.: Gas-phase ozone oxidation of monoterpenes: Gaseous and particulate products, *J. Atmos. Chem.*, 34, 207–258, 1999a.
- Yu, J. Z., Griffin, R. J., Cocker, D. R., Flagan, R. C., Seinfeld, J. H., and Blanchard, P.: Obser-

ACPD

7, 18015–18042, 2007

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

vation of gaseous and particulate products of monoterpene oxidation in forest atmospheres, Geophys. Res. Lett., 26, 1145–1148, 1999b.

Zheng, M., Cass, G. R., Schauer, J. J., and Edgerton, E. S.: Source apportionment of PM_{2.5} in the southeastern United States using solvent-extractable organic compounds as tracers,

Environ. Sci. Technol., 36, 2361–2371, 2002.

Zheng, M., Ke, L., Edgerton, E. S., Schauer, J. J., Dong, M. Y., and Russell, A. G.: Spatial distribution of carbonaceous aerosol in the southeastern United States using molecular markers and carbon isotope data, J. Geophys. Res.-Atmos., 111, doi:10.1029/2005JD006777 2006.

Zheng, M., Cass, G. R., Ke, L., Wang, F., Schauer, J. J., Edgerton, E. S., and Russell, A. G.: Source apportionment of daily fine particulate matter at Jefferson street, Atlanta, GA, during summer and winter, J. Air Waste Manage., 57, 228–242, 2007.

Zielinska, B., Sagebiel, J., McDonald, J. D., Whitney, K., and Lawson, D. R.: Emission rates and comparative chemical composition from selected in-use diesel and gasoline-fueled vehicles, J. Air Waste Manage., 54, 1138–1150, 2004.

ACPD

7, 18015–18042, 2007

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Table 1. Ratios of major resin acids to levoglucosan in source emissions and ambient samples.

Compounds	Softwood Emissions					Ambient Samples		
	Loblolly Pine ^a	Slash Pine ^a	White pine ^b	Hemlock ^b	Balsam fir ^b	Before_Fire	Smoke	After_Fire
Pimaric acid	0.069	0.029	0.008	0.001	0.001	0.001	0.000	0.002
Sandaracopimaric acid	0.013	0.011	0.026	0.002	0.001	0.003	0.001	0.011
Dehydroabietic acid	0.339	0.141	0.149	0.017	0.028	0.041	0.034	0.248
Abietic acid	0.801	0.056	0.391	0.021	0.240	0.001	0.000	0.002
Abieta-6,8,11,13,15-pentae-18-oic acid	0.007	0.003	0.005	0.000	0.002	0.000	0.000	0.000
Abieta-8,11,13,15-tetraen-18-oic acid	0.021	0.000	0.010	0.001	0.003	0.001	0.001	0.001
7-Oxodehydroabietic acid	0.009	0.004	0.005	0.001	0.000	0.007	0.015	0.042

^a Softwood species tested by Fine et al. (Fine et al., 2002);

^b Softwood species tested by Fine et al. (Fine et al., 2001).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Source composition profiles from the aged plume on the event and the previous prescribed burning emission ($\text{ng}/\mu\text{g PM}_{2.5}$).

Compound	Aged Plume ^a	PPM _{TC} ^b	Lee et al. ^c	Compound	Aged Plume ^a	PPM _{TC} ^b	Lee et al. ^c
	Fraction ^d ±std ^e	Fraction±std	Fraction±std		Fraction±std	Fraction±std	Fraction±std
<i>n</i> -alkanes							
Tetracosane	0.0375±0.0208	0.0508±0.0284	0.0836±0.0528	Triacotane	0.0451±0.0116	0.0610±0.0164	0.1168±0.0715
Pentacosane	0.0706±0.0294	0.0956±0.0404	0.1506±0.0954	Hentriacotane	0.2561±0.0731	0.3467±0.1020	0.1737±0.1054
Hexacosane	0.0225±0.0159	0.0305±0.0217	0.1374±0.0913	Dotriacotane	0.0292±0.0075	0.0395±0.0106	0.0000±0.0001
Heptacosane	0.1728±0.0497	0.2339±0.0693	0.1505±0.1137	Tritriacotane	0.0527±0.0156	0.0713±0.0217	0.0542±0.0480
Octacosane	0.0470±0.0171	0.0637±0.0236	0.0695±0.0416	Tetrtiacotane	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001
Nonacosane	0.3115±0.0881	0.4217±0.1231	0.4860±0.2950				
Branch-alkanes							
iso-Nonacosane	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001	iso-Hentriacotane	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001
anteiso-Triacotane	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001				
Hopanes							
17 α (H)-21 β (H)-29-Norhopane	0.0022±0.0051	0.0029±0.0069	0.0000±0.0001	22R,17 α (H),21 β (H)-Homohopane	0.0018±0.0021	0.0025±0.0029	0.0000±0.0001
17 α (H)-21 β (H)-Hopane	0.0016±0.0047	0.0021±0.0064	0.0000±0.0001	22S,17 α (H),21 β (H)-Bishomohopane	0.0009±0.0013	0.0012±0.0017	0.0000±0.0001
22,29,30-Trisnorhomohopane	0.0022±0.0020	0.0029±0.0028	0.0000±0.0001	22R,17 α (H),21 β (H)-Bishomohopane	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001
22,29,30-Trisnorhopane	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001	22S,17 α (H),21 β (H)-Trishomohopane	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001
22S,17 α (H),21 β (H)-Homohopane	0.0028±0.0028	0.0038±0.0038	0.0000±0.0001	22R,17 α (H),21 β (H)-Trishomohopane	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001
Steranes							
20S,R-5 α (H),14 β (H),17 β (H)-Cholestanes	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001	20S,R-5 α (H),14 β (H),17 β (H)-Ergostanes	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001
20R-5 α (H),14 α (H),17 α (H)-Cholestane	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001	20S,R-5 α (H),14 β (H),17 β (H)-Sitostanes	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001
PAHs							
Fluoranthene	0.0000±0.0001	0.0000±0.0001	0.0539±0.0310	Benzo(j)fluoranthene	0.0004±0.0002	0.0006±0.0003	0.0077±0.0049
Acenaphanthrylene	0.0009±0.0004	0.0012±0.0005	0.0168±0.0105	Benzo(e)pyrene	0.0039±0.0024	0.0053±0.0033	0.0300±0.0168
Pyrene	0.0000±0.0001	0.0000±0.0001	0.0643±0.0363	Benzo(a)pyrene	0.0006±0.0016	0.0008±0.0022	0.0179±0.0100
Retene	0.0284±0.0077	0.0385±0.0108	0.2102±0.1267	Perylene	0.0000±0.0001	0.0000±0.0001	0.0023±0.0015
Benzo(ghi)fluoranthene	0.0018±0.0018	0.0025±0.0025	0.0585±0.0517	Indeno(1,2,3-cd)fluoranthene	0.0024±0.0011	0.0033±0.0015	0.0000±0.0001
Cyclopenta(cd)pyrene	0.0000±0.0001	0.0000±0.0001	0.1740±0.1058	Indeno(1,2,3-cd)pyrene	0.0028±0.0025	0.0038±0.0034	0.0444±0.0265
Benzo(a)anthracene	0.0006±0.0017	0.0008±0.0023	0.0482±0.0268	Picene	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001
Chrysene/Triphenylene	0.0071±0.0041	0.0097±0.0056	0.0589±0.0332	Benzo(ghi)perylene	0.0026±0.0041	0.0036±0.0056	0.0215±0.0120

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. (Continued.)

Compound	Aged Plume ^a	PPM _{TC} ^b	Lee et al. ^c	Compound	Aged Plume ^a	PPM _{TC} ^b	Lee et al. ^c
	Fraction ^d ± std ^o	Fraction ± std	Fraction ± std		Fraction ± std	Fraction ± std	Fraction ± std
PAHs							
Benzo(b)fluoranthene	0.0043±0.0028	0.0059±0.0039	0.0304±0.0170	Coronene	0.0039±0.0037	0.0053±0.0051	0.0044±0.0031
Benzo(k)fluoranthene	0.0028±0.0019	0.0038±0.0026	0.0245±0.0136				
Resin Acids							
8,15-Pimaredienoic acid	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001	Abietic acid	0.0112±0.0043	0.0152±0.0059	0.0000±0.0001
Pimaric acid	0.0185±0.0055	0.0251±0.0077	1.4856±0.8256	Abieta-6,8,11,13,15-pentae-18-oic acid	0.0088±0.0025	0.0120 ± 0.0036	0.0000±0.0001
Sandaracopimaric acid	0.0558±0.0169	0.0756±0.0235	0.0000±0.0001	Abieta-8,11,13,15-tetraen-18-oic acid	0.0263±0.0072	0.0356±0.0101	0.0000±0.0001
Isopimaric acid	0.0000±0.0001	0.0000±0.0001	1.7830±0.9994	7-Oxodehydroabietic acid	0.7430±0.1978	1.0058±0.2773	0.0000±0.0001
Dehydroabietic acid	1.5456±0.4331	2.0923±0.6051	20.073±11.450				
Aromatic Acids							
1,2-Benzenedicarboxylic acid	0.1185±0.0359	0.1604±0.0499	0.0088±0.0062	1,3-Benzenedicarboxylic acid	0.0286±0.0112	0.0387±0.0154	0.0000±0.0001
1,4-Benzenedicarboxylic acid	0.0436±0.0207	0.0591±0.0284	0.0054±0.0047				
Alkanoic Acids							
Tetradecanoic acid	0.0741±0.0248	0.1004±0.0343	1.2089±0.7432	Tricosanoic acid	0.1450±0.0392	0.1963±0.0548	0.3941±0.2273
Pentadecanoic acid	0.0359±0.0135	0.0486±0.0187	0.3838±0.2446	Tetracosanoic acid	0.5537±0.1509	0.7496±0.2112	2.9835±1.7399
Hexadecanoic acid	0.2784±0.1320	0.3769±0.1808	4.0548±2.3375	Pentacosanoic acid	0.0488±0.0133	0.0660±0.0186	0.2308±0.1332
Heptadecanoic acid	0.0207±0.0077	0.0280±0.0106	0.1494±0.0907	Hexacosanoic acid	0.3011±0.0811	0.4077±0.1136	2.2120±1.2819
Octadecanoic acid	0.1343±0.0642	0.1818±0.0879	1.3508±0.8111	Heptacosanoic acid	0.0172±0.0046	0.0233±0.0064	0.0850±0.0506
Nonadecanoic acid	0.0262±0.0075	0.0355±0.0105	0.2941±0.1870	Octacosanoic acid	0.1031±0.0276	0.1395±0.0387	0.5953±0.3501
Eicosanoic acid	0.1761±0.0487	0.2363±0.0681	0.7198±0.4371	Nonacosanoic acid	0.0216±0.0056	0.0292±0.0078	0.1014±0.0647
Heneicosanoic acid	0.0740±0.0202	0.1002±0.0283	0.1739±0.1009	Triacontanoic acid	0.1119±0.0299	0.1516±0.0419	0.4837±0.3137
Docosanoic acid	0.3794±0.1031	0.5136±0.1444	1.0757±0.6382				
Alkenoic Acids							
9-Hexadecenoic acid	0.0000±0.0001	0.0000±0.0001	0.3058±0.2272	9-Octadecenoic acid	0.0000±0.0001	0.0000±0.0001	1.4690±0.8808
9,12-Octadecanedienoic acid	0.0000±0.0001	0.0000±0.0001	1.3574±0.9208				
Alkanedioic Acids							
Propanedioic acid	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001	Hexanedioic acid	0.0882±0.0264	0.1195±0.0367	0.0159±0.0132
Methylpropanedioic acid	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001	Heptanedioic acid	0.0000±0.0001	0.0000±0.0001	0.0180±0.0159
Butanedioic acid	0.8009±0.2267	1.0842±0.3165	0.2388±0.1533	Octanedioic acid	0.1487±0.0409	0.2013±0.0572	0.0358±0.0280

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. (Continued).

Compound	Aged Plume ^a	PPM _{fc} ^b	Lee et al. ^c	Compound	Aged Plume ^a	PPM _{fc} ^b	Lee et al. ^c
	Fraction ^d ±std ^e	Fraction±std	Fraction±std		Fraction±std	Fraction±std	Fraction±std
Alkanedioic Acids							
Methylbutanedioic acid	0.1716±0.0470	0.2323±0.0658	0.0000±0.0001	Nonanedioic acid	0.2986±0.0835	0.4043±0.1167	0.1558±0.1006
Pentanedioic acid	0.1198±0.0347	0.1622±0.0484	0.0489±0.0317				
Others							
Nonanal	0.0125±0.0088	0.0170±0.0120	0.0000±0.0001	Benz(de)anthracen-7-one	0.0036±0.0023	0.0049±0.0032	0.0000±0.0001
Sinapyl aldehyde	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001	3,5-Dimethoxy-4-hydroxyacetophenone	0.0275±0.0071	0.0372±0.0100	0.0000±0.0001
Acetonylsyringol	0.0581±0.0181	0.0786±0.0252	0.0000±0.0001	Levogluconan	45.911±12.692	62.151±17.746	57.087±32.377
Coniferyl aldehyde	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001	Cholesterol	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001
Propionylsyringol	0.0000±0.0001	0.0000±0.0001	0.0000±0.0001				
OC	593.85±180.15	450.19±250.55	602.50±185.20	Cl ⁻	0.4355±0.0898	0.5895±0.1287	5.2700±2.8900
EC	22.962±12.421	31.085±16.961	39.200±11.300	NH ₄ ⁺	13.119±4.4281	17.760±6.1272	1.0700±1.0800
Al	0.0000±0.0001	0.0000±0.0001	0.2290±0.4260	NO ₃ ⁻	19.310±5.1844	26.141±7.2624	4.4000±2.9900
Si	0.2889±0.2276	0.3911±0.3094	0.1860±0.2580	SO ₄ ²⁻	13.800±8.5722	18.682±11.680	2.4500±1.1200
K ⁺	2.2261±0.5044	3.0135±0.7159	6.4900±4.3500				

^a source composition profiles where individual chemical species are normalized to the fire-caused total PM_{2.5} mass on the event day;

^b source composition profiles where individual chemical species are normalized to the fire-caused primary PM_{2.5} mass estimated on the event day;

^c source composition profiles from the prescribed burning emission in Georgia (Lee et al., 2005);

^d fraction of chemical species in the associated PM_{2.5} mass (ng/μg PM_{2.5});

^e standard deviation.

Carbonaceous aerosols in an aged prescribed fire plume

B. Yan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Carbonaceous
aerosols in an aged
prescribed fire plume**

B. Yan et al.




Fig. 1. 24-h average ambient concentrations of OC and levoglucosan observed before, during and after the event day at Atlanta area, GA.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Carbonaceous
aerosols in an aged
prescribed fire plume**

B. Yan et al.




Fig. 2. 24-h average concentrations of resin acids and PAHs observed before, during and after the event day at Atlanta area, GA.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Carbonaceous
aerosols in an aged
prescribed fire plume**

B. Yan et al.




Fig. 3. 24-h average concentrations of *n*-alkanes and *n*-alkanoic acids observed before, during and after the event day at Atlanta area, GA.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Carbonaceous
aerosols in an aged
prescribed fire plume**

B. Yan et al.




Fig. 4. 24-h average concentrations of alkanedioic acids, hopanes and steranes observed before, during and after the event day at Atlanta area, GA.
18040

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Carbonaceous
aerosols in an aged
prescribed fire plume**

B. Yan et al.




Fig. 5. 24-h average concentrations of other compounds (cholesterol, 2-methyltetrols, pinonic acid and pinic acid) observed before, during and after the event day at Atlanta area, GA.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Carbonaceous
aerosols in an aged
prescribed fire plume**

B. Yan et al.




Fig. 6. Comparison between source composition profiles developed from the aged biomass burning plume on the event and the prescribed burning emission measured by Lee et al. (2005). The profiles are composed of individual species fractions, normalized to primary PM_{2.5} mass. A log-log scale is used.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion